

Review Article

Factors in fresh market stone fruit quality

Carlos H. Crisosto, F. Gordon Mitchell and Scott Johnson
Pomology Department, University of California, Davis
Kearney Agricultural Center, 9420 South Riverbend Avenue, Parlier,
CA 93648, USA

ABSTRACT

Postharvest quality in stone fruits is ultimately defined in terms of consumer satisfaction, and includes appearance, texture and flavor, and nutritional value and safety. These result from all decisions and practices beginning with site and cultivar selection, and extending through cultural practices and all stages of postharvest handling. The development of high quality fruit is influenced by such cultural practices as mineral nutrition, irrigation, tree training, pruning and fruit thinning. Harvest maturity is of special importance because of the detrimental effects on quality of harvesting at either too low or too high a maturity. Fruit deterioration, from such factors as fruit rotting organisms, water loss, mechanical bruising and physiological disorders, will adversely impact quality in the market. Of special concern is the need to rapidly cool stone fruits and maintain them near 0C throughout marketing. Only by attention to all aspects of production and postharvest handling can high quality be achieved and protected in fresh stone fruits.

INTRODUCTION

In recent years the production and fresh marketing of stone fruits (apricots, cherries, nectarines, peaches and plums) has increased rapidly (LaRue and Johnson, 1989). This has been due in large part to successes in achieving long distance transport to receiving markets. While some movement of southern hemisphere stone fruits to northern hemisphere markets has occurred for many years, only in recent years have these fruits become available worldwide through much of the year. The marketing successes have resulted from a combination of new cultivars and improvements in fruit handling and distribution.

If the volume of shipments is to continue or increase, greater attention must be given to the production and protection of high quality stone fruits. The California industry has seen increasing demand in the U.S. for high maturity fresh stone fruit. Special packing has been developed to supply this premium market, but the handling requirements limit the volume available. This has caused both the industry and University researchers to evaluate the potential for improving the quality (and hopefully advancing the maturity) of fruits packed in large volume operations. To achieve this, all aspects of preharvest and postharvest factors affecting fruit quality must be explored. This paper reviews some of the recent work in California.

EFFECTS OF CULTURAL PRACTICES ON FRUIT QUALITY

While postharvest horticulturists have generally focused on harvest maturity (Kader and Mitchell, 1989) and temperature management (Mitchell, 1987; 1989) as their starting point, many earlier decisions will profoundly influence the postharvest performance of the fruit. Some of these decisions even predate orchard planting, such as location (climate), site (soil) and cultivar selection. A number of preharvest cultural practices can influence postharvest quality and performance. There are also many handling practices during harvest, packing and distribution that affect quality, as well as the deleterious effect of various fruit deterioration problems.

The studies and observations discussed here relate mainly to stone fruit production in California. The central valley of California, where most stone fruits are grown, is characterized by a Mediterranean climate, with hot, dry summers. Stone fruits are grown under summer irrigation, primarily in deep, alluvial soils.

Mineral Nutrition

Research over the last 10 years has established that peaches and nectarines should be kept between 2.6 and 3.0% leaf nitrogen (N) for

best fruit quality. N deficiency leads to small fruit with poor flavor and unproductive trees. Conversely, many negative effects from excessive N levels have demonstrated (Daane *et al.*, 1994; Michailides *et al.*, 1992).

Response of peach and nectarine trees to N fertilization is dramatic; high N levels stimulate vigorous vegetative growth, shading out of lower fruiting wood, and adversely affecting fruit quality. Fruit maturity is delayed, percent red coloration is decreased, and fruit size is not increased as compared to optimum N levels (Daane *et al.*, 1994). Heavily fertilized trees have also been observed to be more susceptible to brown rot (*Monilinia fructicola*) and various insects and mites. Although high N trees may look healthy and lush, excess N does not increase fruit size, production or soluble solids concentration (SSC).

Current research is evaluating sources of N other than commercial fertilizers. Preliminary results indicate that some materials which increase soil organic matter (composts, manures, cover crops, etc.) may improve some aspects of fruit quality, such as SSC, firmness and insect and disease resistance.

Foliar sprays

We have little detailed research on foliar sprays which supply small amounts of mineral nutrients. However, our observations and reports in the literature suggest that these sprays have little effect on fruit growth or quality. Recent research (Cheng and Crisosto, 1994; Crisosto *et al.*, 1993b) suggests that these sprays should be treated with caution, since their heavy metal content (Fe, Al, Cu, etc.) may contribute to fruit skin discoloration.

Our recent work with three commercial Ca foliar sprays on peach and nectarine (applied every 14 days, starting 2 weeks after full bloom and continuing until one week before harvest) showed no reduction in skin discoloration or increase in fruit quality of mid- or late-season cultivars (Crisosto *et al.*, 1993c). Also, these foliar sprays did not affect fruit SSC or firmness, or Ca content of leaves or fruit flesh at harvest or after 2 or 4 weeks in cold storage.

Irrigation

Trees supplied with optimum amounts of water (100% evapotranspiration, E.T.) during the season will produce maximum fruit size (Crisosto *et al.*, 1994). However, higher SSC levels can be obtained by imposing moderate water stress during fruit growth prior to harvest. This will also reduce fruit size, but in some cases only slightly. When 'O'Henry' peach were held under moderate water stress (50% E.T. in the month prior to harvest), the stored fruit did not develop dry and mealy texture (internal breakdown symptoms). Moderate water stress can also reduce vegetative growth, thus maintaining lower fruiting wood, especially in high density plantings. Over-irrigation (150% E.T. in the month prior to harvest) can be detrimental to long term productivity. Although fruit size may increase in the short term, the resulting excess vegetative growth can lead to shading out of lower fruiting wood and loss of yield. Saturated soil conditions can lead to tree decline.

In earlier work, extreme water stress resulted in small astringent fruit and unproductive trees. Imposing severe water stress in late summer (no irrigation during the growing season) on an early maturing cultivar decreased fruit quality the following year by increasing doubled and deep-sutured fruit.

Canopy position

During the last three seasons, we have evaluated fruit quality and postharvest performance of several peach, nectarine, and plum cultivars (Day *et al.*, 1992; Mitchell *et al.*, 1990; Saenz, 1991). Large differences in SSC, acidity, fruit size and storage potential were detected between fruit taken from the top, outside and inside canopy positions of open-vase trained trees. These differences in postharvest characteristics related to canopy position, became smaller when trees of early cultivars were trained to a V-shape, with the V oriented across the direction of the rows.

Thinning

Fruitlet thinning will increase fruit size while also reducing total yield, and thus a balance between yield and fruit size must be achieved (Day *et al.*, 1992). Generally, maximum profit does not occur at maximum marketable yield since larger fruit bring a higher market price. Leaving too many fruit on a tree not only reduces fruit size but also decreases their SSC. Thus fruit quality can be sacrificed in several ways by incorrect thinning. Grower experience is the best determinant of the optimum thinning level for each orchard and cultivar.

Pruning

The greater the light interception by an individual fruit and its surrounding leaves the better its quality (including fruit color, size, SSC, and flavor). Fruit in the top of the tree, for example, always have better quality than fruit in the lower, shaded part of the canopy (Day *et al.*, 1992). The differences can sometimes be substantial, even though the lower fruit remain on the tree for a longer period to reach maturity.

Prudent summer pruning practices which increase light penetration into the canopy will generally improve fruit quality. For example, the removal of interior water sprouts can significantly increase light penetration and improve fruit size, color and SSC of lower position fruit. However extensive summer pruning which removes many leaves surrounding the fruit can have the opposite effect, since these leaves supply carbohydrates to the fruit. In the same way, the practice of removing ("pulling") leaves around fruit to increase fruit color may decrease fruit size and SSC.

POSTHARVEST FACTORS AFFECTING STONE FRUIT QUALITY

These stone fruits as a group typically have a relatively short postharvest life (Mitchell and Kader, 1989a). They can quickly pass from ideal ripeness to over-maturity, depending in part on temperature and handling exposures (Kader and Mitchell, 1989). They will readily lose water and shrivel (with stem browning in

cherries), as a result of movement of water vapor from the fruit to the environment, depending upon surface characteristics of the fruit, relative humidity, air velocity and temperature (Crisosto *et al.*, 1993a; Mitchell, 1987; 1989). They can be quickly destroyed by fruit rotting organisms, depending upon the level of inoculum, surface wounds from handling, fungicide treatment and temperature. All except cherry can also become unmarketable as a result of internal breakdown, a chilling injury problem that develops in many cultivars, depending in part upon harvest maturity, temperature and holding duration.

Harvest maturity

The maturity of stone fruits at harvest will determine their ability to achieve high eating quality, their susceptibility to mechanical injuries, their postharvest performance and their potential postharvest life. Any maturity index should clearly separate fruit based on physiological maturity, and any legal standard should be independent of growing conditions or location (Crisosto, 1992; Kader and Mitchell, 1989).

Fruit harvested at too high a maturity will be incapable of withstanding the rigors of postharvest handling and distribution, may have increased susceptibility to invasion by fruit rotting organisms (Mitchell *et al.*, 1991). These fruit will have a short postharvest life, and may develop undesirable off flavors and mealy texture.

Fruit harvested at too low a maturity will be incapable of ripening to their potential flavor and texture qualities. They will also lose water more readily, and may be at increased risk of physiological deterioration, especially if susceptible to internal breakdown.

With the increasing volume of fresh stone fruits entering long distance marketing, there has been a tendency to harvest the fruit at lower maturity; the idea being that they would be better able to carry to distant markets. The result has often been negative, however, because such fruit are more subject to shrivel and to the development of internal breakdown symptoms.

Handling injuries

Mechanical injuries can occur at any time from harvest through packing, handling and transportation (Mitchell and Kader, 1989b). Impact and compression bruises are normally the result of rough handling somewhere in the system. They typically follow abuses during picking, filling or transfer of field containers, packing line operations, transport and distribution. Recognizing that firmer fruit will withstand greater abuse, producers then tend to harvest at lower maturity. Our studies have shown that careful monitoring and supervision throughout the system can reduce the injury potential. This includes training and supervision of pickers, field handlers, packing line workers, assemblers and loaders. Recently developed instrumentation can be useful in identifying potential injury locations on packing lines and other equipment.

In recent studies (Mitchell *et al.*, 1989; Mitchell *et al.*, 1990; Mitchell *et al.*, 1991) the injury susceptibility of various nectarine, peach and plum cultivars increased at flesh firmness below about 3.5 kg-force. At similar firmness, plums were more resistant to bruising than were nectarines or peaches. Because of their greater susceptibility and their normal flesh firmness variability at harvest, nectarines and peaches that pass over typical packing lines in California should probably have an average flesh firmness of at least 4.5 to 5.5 kg-force. It would be desirable to improve the performance of these packing lines so harvest maturity could be advanced.

A limited U.S. market has recently developed for higher maturity fruit. Called "Tree Ripe", such fruit are usually just harvested at a higher than normal maturity (Mitchell *et al.*, 1989; Mitchell *et al.*, 1990). Because they will not withstand the rigors of typical commercial packing lines, these softer fruit are packed on small, labor intensive hand packing lines. While the demand for such fruit is strong, the volume is limited by availability of suitable packing facilities and labor.

Vibration or abrasion bruising can result from fruit movement or rubbing during handling or transportation (Mitchell and Kader, 1989a). This injury can cause discoloration of the fruit surface, accelerate water loss, and inflict fresh wounds for pathogen penetration. Incidence can be reduced by avoiding opportunities for fruit to abrade during handling, and by packing the fruit so they will be immobilized during transport.

Recent studies of "black-staining", of peaches and nectarines (Cheng and Crisosto, 1994; Crisosto *et al.*, 1993b) have shown that discoloration can be the result of metal ion contamination of wound areas, especially by Fe, Al and Cu ions. Even foliar nutrient sprays and certain orchard fungicides can cause a problem if applied too close to harvest. Brushing and waxing of these fruits may increase susceptibility to the disorder.

Internal breakdown (chilling injury)

This disorder has not been reported on cherries (Mitchell and Kader, 1989a), but occurs on many cultivars of all other stone fruits (Table 1). The injury results from prolonged exposure to temperatures below 10°C, and develops most quickly and severely in fruit held at 5C (Table 2).

Table 1. Postharvest performance rating of different stone fruit cultivars grown in California (Modified and updated from Mitchell and Kader, 1989).

Cultivar	Internal Breakdown @5°C	MarketLife Wks @0°C	Cultivar	InternalBreakdown @5°C	MarketLife Wks @0°C
Nectarine					
August Red	M	3	Mayfire	L	6
Autumn Grand	H	2-3	May Grand	L	6
Fairlane	H	3	Red Free	L	6
Fantasia	M	4-6	Red Grand	M	3-5
Flamekist	M	3	Royal Giant	M	4
Flaming Red	M	3	September Grand	M	3-4
Flavortop	L	4-5	September Re	M	3
Independence	L	5-6	Sparkling Red	M	5
July Red	L	6	Spring Red	L	6
Juneglo	L	6	Summer Grand	L	5
May Diamond	L	6	Summer Red	M	5
Mavglo	L	4			
Peach					
Autumn Gem	H	1	Lacey	M	6
Belmont	H	1	Merrill Gemfree	L	6
Cal Red	H	2	O'Henry	M	4
Carnival	H	2	Parade	H	3
Cassie	H	2	Red Cal	H	2
Early Fairtime	H	2	Redtop	L	4
Early O'Henry	H	4	Regina	L	3
Elegant Lady	H	4	Sparkle	M	3
Fairtime	H	2	Springcrest	H	3-4
Fire Red	H	2	Springgold	H	3
Flamecrest	L	3	Spring Lady	L	5
Flavorcrest	H	4	Summer Lady	H	5
June Lady	M	3	Suncrest	M	2
Kings Lady	H	2	Windsor	H	2
Plum					
Ambra	H	2	Moyer	L	5
Angeleno	M	8-10	Nubiana	H	2
Black Beaut	L	3	President	L	3
Casselman	M	5-6	Queen Ann	M	4
Catalina	L	6	Queen Rosa	M	4
El Dorado	M	3-5	Red Beaut	H	1-2
French Prune	L	>8	Red Rosa	H	4
Friar	M	3-4	Rosemary	L	4
Frontier	H	>4	Royal Diamond	M	6
Grand Rosa	H	3	Roysum	M	4
July Santa Rosa	H	3	Santa Rosa	L	3-5
Kelsey	H	2	Simka	M	3
Laroda	L	4	Spring Beaut	L	2
Late Santa Rosa	L	3	Wickson	L	4

Table 2. Percent of marketable stone fruit following 2 different temperature regimes.

Cultivar	Percent Juicy Fruit	
	0°C for 6 weeks	5°C for 3 weeks
Peaches		
Spring Lady	80	17
Springcrest	95	35
Flavorcrest	85	42
Redtop	42	27
O'Henry	67	14
Fairtime	67	23
Nectarines		
Juneglo	100	100
Sparkling June	86	100
Fantasia	91	40
September Red	49	13
Fairlane	50	9

Crisosto unpublished data.

Most susceptible cultivars require 2 to 3 weeks exposure below 10°C for symptoms to occur, but a few peach cultivars will show problems after only one week of exposure. One or more visual symptoms can appear, including dry or mealy texture, failure to ripen, flesh browning, reddening or translucency, and dark discoloration of the stone cavity; and these symptoms are always accompanied by a loss of characteristic flavor of the fruit. Various conditioning treatments have been tried to delay development, but none are currently commercialized (Nanos and Mitchell 1991a; 1991b). While the disorder will develop in fruit held at or below 0°C, symptoms are less severe, and development is delayed. Thus, it is important that those cultivars be promptly cooled and held at temperatures of 0°C or below (depending on the freezing point of the fruit).

Temperature management

A summary of the effect of temperature on stone fruits is presented in Table 3. Temperature effects depend on the duration of exposure to a particular temperature and to the postharvest age of the fruit (Mitchell 1987; 1989). Stone fruits as a group are very responsive to high temperature exposure, such as delays between harvest and cooling. Problems can include high temperature injury, rapid softening, excess water loss and shrivel, and greater sensitivity to other disorders. Thus protection from heating after harvest and rapid movement to cooling are important. This is especially important with high maturity fruit, to avoid excessive flesh softening, but can also help to reduce the detrimental effects of internal breakdown.

Sweet cherries should ideally be cooled to near 0°C within 3-4 hours of harvest (Crisosto *et al.*, 1993a). Plums and normal maturity peaches and nectarines should be cooled to 5-10°C within 6-8 hours, and to 0°C within 24 hours of harvest; high maturity peaches and nectarines perform best if cooled to near 0°C within about 6-8 hours of harvest.

Stone fruits should be held between -0.5 and 0°C throughout the postharvest period, unless special conditioning treatments are being

applied. Care should be taken to avoid prolonged exposure to intermediate temperatures because of the potential for flesh softening, shrivel, fruit rot development and internal breakdown (Mitchell and Kader, 1989a).

Table 3. Temperature effects on stone fruits.

Temperature (°C)	Fruit response
>40	Skin scald and flesh breakdown
37 to 40	Failure to ripen
28 to 36	Abnormal ripening (mealiness, off-flavors)
15 to 27	Normal ripening (higher temperatures induce faster ripening)
0 to 10	Chilling injury/internal breakdown (severity depends on species and cultivar)
2 to 8	Most severe chilling injury
-0.5 to 1.5	Minimum flesh softening
-0.5 to 0	Best storage life
<-0.8	Possible freezing (depending on soluble solids concentration)

Fruit of cultivars that are susceptible to internal breakdown require a period of exposure, usually 2 weeks or more, to develop symptoms, and temporary holding near 5°C will not cause symptoms to develop.

Thus prompt cooling and overnight holding at temperatures between 5 and 10°C to await next-day packing should pose no problem, provided the fruit are promptly cooled and held near 0°C following packing. If packing is to be delayed beyond the next day, then fruit awaiting packing should be cooled to near 0°C. Unfortunately, undesirable holding temperatures (around 4-6°C) that are often encountered inside loads of these fruits during transport to consumer markets can exaggerate deterioration.

FINAL COMMENTS

The production and marketing of high quality stone fruits requires attention to all aspects of the system, from site and cultivar selection, through all cultural practices during fruit production, to harvest maturity, fruit selection and field handling, grading and packing, temperature management and distribution. Just as most cultivars of the past are unsuited to today's markets, so most of the older handling practices are not adequate to protect the fruit during long distance marketing. Perhaps this is most notable in temperature management, where newer methods of rapid cooling and low temperature holding are essential.

REFERENCES

- Cheng, G.W.; Crisosto, C.H.. (1994) Development of dark skin discoloration on peach and nectarine fruit in response to exogenous contaminations. *Journal of the American Society for Horticultural Science* **119**, 529-533.
- Crisosto, C. H. (1992) Stone fruit maturity indices: a descriptive review. In *1992 Research Reports for California Peaches and Nectarines, California Tree Fruit Agreement*. Sacramento, California, USA. 10 pp.

- Crisosto, C.H.; Garner, D.; Doyle, J.; Day, K. R.. (1993a) Relationship between fruit respiration, bruising susceptibility, and temperatures in sweet cherries. *HortScience* **28**, 132-135.
- Crisosto, C. H.; Johnson, R. S.; Luza, J. G.; Crisosto, G. M.. (1994) Irrigation regimes affect fruit soluble solids content and the rate of water loss of 'O'Henry' peaches. *HortScience* **29**, 1169-1171.
- Crisosto, C.H.; Johnson, R.S.; Luza, J.; Day, K.R. (1993) Incidence of physical damage on peach and nectarine skin discoloration development: anatomical studies. *Journal of the American Society for Horticultural Science* **118**, 796-800.
- Crisosto, C.H.; Johnson, R.S.; Cheng, G.W.; Day, K.R, Garner, D.; Crisosto, G. (1993) Skin discoloration on peach and nectarine fruit. In *1993 Research Reports for California Peaches and Nectarines, California Tree Fruit Agreement*. Sacramento, California, USA. 15pp.
- Daane, K.M.; Johnson, R.S.; Michailides, T.J.; Crisosto, C.H.; Dlott, J.W.; Ramirez, H.T.; Yokota, G.T.; Morgan, D.P. (1994) Nitrogen fertilization affects nectarines fruit yield, storage qualities, and susceptibility to brown rot and insect damage. *California Agriculture* (In press).
- Day, K.R.; Johnson, R.S.; DeJong, T.M.; Crisosto, C.H. (1992) Comparison of high density training systems and summer pruning techniques and timing. In *1992 Research Reports for California Peaches and Nectarines, California Tree Fruit Agreement*. Sacramento, California, USA. 4pp.
- Kader, A.A.; Mitchell, F.G. (1989) Maturity and quality. In LaRue, J.H.; Johnson, R.S. (eds) *Peaches, Plums and Nectarines - Growing and Handling for Fresh Market*. University of California Division of Agriculture and Natural Resources, Publication No. 3331. pp. 191-196.
- LaRue, J. H. and Johnson, R. S. (eds.). (1989) *Peaches, Plums and Nectarines - Growing and Handling for Fresh Market*. University of California Division of Agriculture and Natural Resources, Publication No. 3331, Oakland, California. 246 pp.
- Michailides, T.J.; Ramirez, H.T.; Morgan, D. P.; Crisosto, C. H.; Johnson, R.S. (1992) Nitrogen fertilization affects fruit quality and susceptibility of nectarine fruits to brown rot. In *1992 Research Reports for California Peaches and Nectarines, California Tree Fruit Agreement*. Sacramento, California, USA. 24 pp.
- Mitchell, F.G. (1987) Influence of cooling and temperature maintenance on the quality of California grown stone fruit. *Revue Internationale du Froid* **10**, 77-81.
- Mitchell, F.G. (1989) Cooling. In LaRue, J.H.; Johnson, R.S. (eds) *Peaches, Plums and Nectarines - Growing and Handling for Fresh Market*. University of California Division of Agriculture and Natural Resources, Publication No. 3331. pp. 209-215.
- Mitchell, F.G.; Kader, A.A. (1989a) Factors affecting deterioration rate. In LaRue, J.H.; Johnson, R.S. (eds) *Peaches, Plums and Nectarines - Growing and Handling for Fresh Market*. University of California Division of Agriculture and Natural Resources, Publication No. 3331. pp. 165-178.
- Mitchell, F.G.; Kader, A.A. (1989b) Storage. In LaRue, J.H.; Johnson, R.S. (eds) *Peaches, Plums and Nectarines - Growing and Handling for Fresh Market*. University of California Division of Agriculture and Natural Resources, Publication No. 3331. pp. 216-222.
- Mitchell, F.G.; Mayer, G.; Biasi, W.; Gulli, D.; Faubion, D. (1989) Selecting and handling high quality stone fruit. In *1989 Research Reports for California Peaches and Nectarines, California Tree Fruit Agreement*. Sacramento, California, USA. 15pp.
- Mitchell, F.G.; Mayer, G.; Johnson, R.S.; Biasi, W. (1990) Factors in selecting and handling high quality stone fruit. In *1990 Research Reports for California Peaches and Nectarines, California Tree Fruit Agreement*. Sacramento, California, USA. 27pp.
- Mitchell, F.G.; Mayer, G.; Saenz, M.; Slaughter, D.; Johnson, R.S.; Biasi, B; Delwiche, M. (1991) Selecting and handling high quality stone fruit for fresh market. In *1991 Research Reports for California Peaches and Nectarines, California Tree Fruit Agreement*. Sacramento, California, USA. 11pp.
- Nanos, G.D.; Mitchell, F.G. (1991a) Carbon dioxide injury and flesh softening following high-temperature conditioning in peaches. *HortScience* **26**, 562-563.
- Nanos, G.D.; Mitchell, F.G. (1991b) High temperature conditioning to delay internal breakdown development in peaches and nectarines. *HortScience* **26**, 882-885.
- Saenz, M.V. (1991) Effect of position in the canopy on the postharvest performance and quality of stone fruit. In *1991 Research Reports for California Peaches and Nectarines, California Tree Fruit Agreement*. Sacramento, California, USA. 19pp.